International
Online Conference



Algebraic and Geometric Methods of Analysis

dedicate to the memory
of Yuriy Trokhymchuk
(17.03.1928-18.12.2019)

May 25-28, 2021
Odesa, Ukraine

НТБ ОНМАКТ

List of topics

- Topological methods in analysis
- Geometric problems of complex and mathematical analysis
- Algebraic methods in geometry
- Differential geometry in the whole
- Geometry and topology of differentiable manifolds
- General and algebraic topology
- Geometric and topological methods in natural sciences

Organizers

- Ministry of Education and Science of Ukraine
- Odesa National Academy of Food Technologies
- Institute of Mathematics of the National Academy of Sciences of Ukraine
- Taras Shevchenko National University of Kyiv
- International Geometry Center
- Kyiv Mathematical Society

Scientific Committee

Drozd Yu.
(*Kyiv, Ukraine*)
Maksymenko S.
(*Kyiv, Ukraine*)
Plaksa S.
(*Kyiv, Ukraine*)
Prishlyak A.
(*Kyiv, Ukraine*)

Bakhtin O.
(*Kyiv, Ukraine*)
Balan V.
(*Bucharest, Romania*)
Banakh T.
(*Lviv, Ukraine*)
Borysenko O.
(*Kharkiv, Ukraine*)

Cherevko Ye.
(*Odesa, Ukraine*)
Fedchenko Yu.
(*Odesa, Ukraine*)
Karlova O.
(*Chernivtsi, Ukraine*)
Kiosak V.
(*Odessa, Ukraine*)
Konovenko N.
(*Odessa, Ukraine*)
Lyubashenko V.
(*Kyiv, Ukraine*)
Matsumoto K.
(*Yamagata, Japan*)
Mormul P.
(*Warsaw, Poland*)

Mykhailyuik V.
(*Chernivtsi, Ukraine*)
Plachta L.
(*Krakov, Poland*)
Pokas S.
(*Odessa, Ukraine*)
Sabitov I.
(*Moscow, Russia*)
Savchenko O.
(*Kherson, Ukraine*)
Sergeeva A.
(*Odesa, Ukraine*)
Shelekhov A.
(*Tver, Russia*)
Zarichnyi M.
(*Lviv, Ukraine*)

НТБ ОНАХТ

Administrative committee

- Egorov B., chairman, rector of the ONAFT;
- Povarova N., deputy chairman, Pro-rector for scientific work of the ONAFT;
- Mardar M., Pro-rector for scientific-pedagogical work and international communications of the ONAFT;
- Fedosov S., Director of the International Cooperation Center of the ONAFT;
- Kotlik S., Director of the P.M. Platonov Educational-scientific institute of computer systems and technologies "Industry 4.0";
- Lishchenko N. Dean of faculty of the computer systems and automation ONAFT

Organizing commitee

Cherevko Ye.
Eftekharinasab K.
Fedchenko Yu.
Feshchenko B.
Khohliyk O.
Klishchuk B.
Konovenko N.
Kravchenko A.
Kuznietsova I.
Maksymenko S.
.
Osadchuk E.
Plakosh A.
Prus A.
Sergeeva A.
Soroka Yu.

НТБ ОНАХТ

Maximal distance minimizers. Examples and properties

Yana Teplitskaya
(SPbSU, Department of Mathematics and Computer Science)
E-mail: janejashka@gmail.com

I will talk about the sets which have the minimal length over the class of closed connected sets $\Sigma \subset \mathbb{R}^2$ satisfying the inequality $\max_{y\in M} \operatorname{dist}(y, \Sigma) \le r$ for a given compact set $M \subset \mathbb{R}^2$ and some given $r > 0$.

For a given compact set $M \subset \mathbb{R}^2$ consider the functional

$$F_M(\Sigma) := \sup_{y\in M} \operatorname{dist}(y, \Sigma),$$

where Σ is a closed subset of $\mathbb{R}^2$ and $\operatorname{dist}(y, \Sigma)$ stands for the Euclidean distance between y and Σ. Consider the class of closed connected sets $\Sigma \subset \mathbb{R}^2$ satisfying $F_M(\Sigma) \le r$ for some $r > 0$. We are interested in the sets of the minimal length (one-dimensional Hausdorff measure) $\mathcal{H}^1(\Sigma)$ over the mentioned class (*minimizers*).

It is known that for all $r > 0$ the set of minimizers is nonempty. It is proven also that for each minimizer of positive length the equality $F_M(\Sigma) = r$ holds. Furthermore the set of minimizers coincides with the set of solutions of the dual problem: minimize F_M over all closed connected sets $\Sigma \subset \mathbb{R}^2$ with prescribed bound on the total length $\mathcal{H}^1(\Sigma) \le l$.

In [1] (for the plane) and in [2] (for the general case) some properties of minimizers have been proven:

(a) A minimizer cannot contain loops (homeomorphic images of circles).

(b) For every point $x \in \Sigma$ one of two statements is true:

  i there exists a point $y \in M$ (may be not unique) such that $\operatorname{dist}(x, y) = r$ and $B_r(y) \cap \Sigma = \emptyset$;

  ii there exists an $\varepsilon > 0$ such that $S_\Sigma \cap B_\varepsilon(x)$ is either a segment or a regular tripod, i.e. the union of three segments with an endpoint in x and relative angles of $2\pi/3$.

The minimizers for some sets M are known (see pictures) although usually this is not an easy task. Recently (see [3]) at the plane the regularity of minimizers was proved.

Theorem 1. *Let Σ be a maximal distance minimizer for a compact set $M \subset \mathbb{R}^2$. Then*

(i) *Σ is a union of a finite number of arcs (injective images of the segment $[0; 1]$).*

(ii) *The angle between each pair of tangent rays at every point of Σ is greater or equal to $2\pi/3$. The number of tangent rays at every point of Σ is not greater than 3. If it is equal to 3, then there exists such a neighbourhood of x that the arcs in it coincide with line segments.*

References

[1] Miranda, Jr., M. and Paolini, E. and Stepanov, E., *On one-dimensional continua uniformly approximating planar sets*. Volume 27 of *Calculus of Variations and Partial Differential Equations*, 2006.

[2] Paolini, E. and Stepanov, E., *Qualitative properties of maximum distance minimizers and average distance minimizers in $\mathbb{R}^n$*, volume 122 of *Journal of Mathematical Sciences*. New York, 2004.

[3] Yana Teplitskaya. *On regularity of maximal distance minimizers*, arXiv preprint arXiv:1910.07630, 2019.



FIGURE 1.1. An example where $M = \partial B_R(O)$, where $R > 4.98r$.

FIGURE 1.2. An example where $M := \partial B_r([AG])$, $\Sigma = [AG]$.



FIGURE 1.3. An example where $M := \{A, B, C\}$, Σ is a tripod.

НТБ ОНАХТ

I. Streltsova *Projective invariants of linear planar 3-webs* **145**

Kh. Sukhorukova *Ultrametric spaces of $*$-measures* **147**

B. M Sultanov *Sweep of surfaces in Galilean space* **148**

Y. Teplitskaya *Maximal distance minimizers. Examples and properties* **149**

M. Tkachuk, S. Plaksa *Analog of Menchov-Trokhimchuk theorem for monogenic functions in three-dimensional commutative algebra* **151**

E. Tosunoğlu, M. E. Kansu *Onto the some dynamic applications via quaternions* **153**

R. M. Tudoran *On dynamical systems with a prescribed globally bp-attracting set* **154**

T. P. Mokritskaya, A. V. Tushev *On certain fractal-based estimations of subsidence volume* **155**

Jun Ueki *Modular knots obey the Chebotarev law* **156**

J. Vielma, A. Guale *The Terence Tao set and the Collatz conjecture* **159**

V. Dilnyi, A. Vinskovska *Conformal mappings in Hardy-type spaces* **160**

I. Vlasenko *On the equivalency classes of weakly conjugated inner mappings* **161**

L. Vyhivska *Extremal problem for non-overlapping domains with free poles* **162**

N. Zorii *Inner harmonic measure for the fractional Laplacian* **163**

П. Задерей, М. Гаєвський *Мультиплікатори в просторах Харді та пов'язаних з ними просторах* **164**

М. І. Піструіл, І. М. Курбатова *Про деякі закономірності квазі-геодезичних відображень узагальнено-рекурентних просторів* **166**

С. М. Покась, И. И. Белокобыльский *Геодезические отображения римановых пространств второго приближения* **167**

НТБ ОНАХТ