International
Online Conference



Algebraic and Geometric Methods of Analysis

dedicate to the memory
of Yuriy Trokhymchuk
(17.03.1928-18.12.2019)

May 25-28, 2021
Odesa, Ukraine

НТБ ОНМАКТ

List of topics

- Topological methods in analysis
- Geometric problems of complex and mathematical analysis
- Algebraic methods in geometry
- Differential geometry in the whole
- Geometry and topology of differentiable manifolds
- General and algebraic topology
- Geometric and topological methods in natural sciences

Organizers

- Ministry of Education and Science of Ukraine
- Odesa National Academy of Food Technologies
- Institute of Mathematics of the National Academy of Sciences of Ukraine
- Taras Shevchenko National University of Kyiv
- International Geometry Center
- Kyiv Mathematical Society

Scientific Committee

Drozd Yu.
(*Kyiv, Ukraine*)
Maksymenko S.
(*Kyiv, Ukraine*)
Plaksa S.
(*Kyiv, Ukraine*)
Prishlyak A.
(*Kyiv, Ukraine*)

Bakhtin O.
(*Kyiv, Ukraine*)
Balan V.
(*Bucharest, Romania*)
Banakh T.
(*Lviv, Ukraine*)
Borysenko O.
(*Kharkiv, Ukraine*)

Cherevko Ye.
(*Odesa, Ukraine*)
Fedchenko Yu.
(*Odesa, Ukraine*)
Karlova O.
(*Chernivtsi, Ukraine*)
Kiosak V.
(*Odessa, Ukraine*)
Konovenko N.
(*Odessa, Ukraine*)
Lyubashenko V.
(*Kyiv, Ukraine*)
Matsumoto K.
(*Yamagata, Japan*)
Mormul P.
(*Warsaw, Poland*)

Mykhailyuik V.
(*Chernivtsi, Ukraine*)
Plachta L.
(*Krakov, Poland*)
Pokas S.
(*Odessa, Ukraine*)
Sabitov I.
(*Moscow, Russia*)
Savchenko O.
(*Kherson, Ukraine*)
Sergeeva A.
(*Odesa, Ukraine*)
Shelekhov A.
(*Tver, Russia*)
Zarichnyi M.
(*Lviv, Ukraine*)

НТБ ОНАХТ

Administrative committee

- Egorov B., chairman, rector of the ONAFT;
- Povarova N., deputy chairman, Pro-rector for scientific work of the ONAFT;
- Mardar M., Pro-rector for scientific-pedagogical work and international communications of the ONAFT;
- Fedosov S., Director of the International Cooperation Center of the ONAFT;
- Kotlik S., Director of the P.M. Platonov Educational-scientific institute of computer systems and technologies "Industry 4.0";
- Lishchenko N. Dean of faculty of the computer systems and automation ONAFT

Organizing commitee

Cherevko Ye.
Eftekharinasab K.
Fedchenko Yu.
Feshchenko B.
Khohliyk O.
Klishchuk B.
Konovenko N.
Kravchenko A.
Kuznietsova I.
Maksymenko S.
.
Osadchuk E.
Plakosh A.
Prus A.
Sergeeva A.
Soroka Yu.

НТБ ОНАХТ

Modular knots obey the Chebotarev law

Jun Ueki
(Tokyo Denki University, 5 Senju Asahi-cho, Adachi-ku, 120-8551, Tokyo, Japan)
E-mail: uekijun46@gmail.com

A *Chebotarev link* in S^3 is an analogue of the set of all prime numbers in $\mathbb{Z}$. It would play an important roll in arithmetic topology, especially when we formulate an analogue of the idelic class field theory for 3-manifolds [Uek21a] (see also [Mor12, Nii14, NU19, Mih19]). Here is the definition:

Definition 1 (The Chebotarev law)**.** Let $(K_i)_i = (K_i)_{i\in\mathbb{N}_{>0}}$ be a sequence of disjoint knots in a 3-manifold M. For each $n \in \mathbb{N}_{>0}$ and $j > n$, put $L_n = \cup_{i\leq n} K_i$ and let $[K_j]$ denote the conjugacy class of K_j in $\pi_1(M - L_n)$. We say that $(K_i)_i$ *obeys the Chebotarev law* if the density equality

$$\lim_{\nu\to\infty} \frac{\#\{n < j \leq \nu \mid \rho([K_j]) = C\}}{\nu} = \frac{\#C}{\#G}$$

holds for any $n \in \mathbb{N}_{>0}$, any surjective homomorphism $\rho : \pi_1(M - L_n) \to G$ to any finite group, and any conjugacy class $C \subset G$.

In order to answer Mazur's question on the existence of such a link in S^3 [Maz12], by using Parry–Pollicott's zeta functions of symbolic dynamics [PP90], McMullen proved a highly interesting theorem:

Proposition 2 ([McM13, Theorem 1.2])**.** *Let $(K_i)_i$ be the closed orbits of any topologically mixing pseudo-Anosov flow on a closed 3-manifold M, ordered by length in a generic metric. Then $(K_i)_i$ obeys the Chebotarev law.*

Applying his theorem to the monodromy suspension flow of the figure-eight knot K and noting that the Chebotarev law persists under Dehn surgeries, he constructed a Chebotarev link containing K in S^3 [McM13, Corollary 1.3]. We refine his construction in two ways to verify the following assertion:

Theorem 3 ([Uek21b, Theorem 3])**.** *Let L be a fibered hyperbolic link in S^3 and let $(K_i)_i$ denote the sequence of knots consisting of the closed orbits of the suspension flow of the monodromy map and L itself. Then $(K_i)_i$ obeys the Chebotarev law, if ordered by length with respect to a generic metric.*

The union $\mathcal{L} = \cup_i K_i$ is a stably Chebotarev link*, that is, for any finite branched cover $h : M \to S^3$ branched along any finite link in $\mathcal{L}$, the inverse image $h^{-1}(\mathcal{L})$ is again Chebotarev.*

One way is to extend McMullen's theorem for *generalized pseudo-Anosov flows*, which allow 1-pronged singular orbits. The other is to invoke the notion of *rational Fried surgeries*, which produce many (generalized) pseudo-Anosov flows.

Our refinement further provides a new example called modular knots, that are also known as Lorenz knots. Let $\mathbb{H}^2 = \{z \in \mathbb{C} \mid \mathrm{Im} z > 0\}$ denote the upper half plane. The unit tangent bundle of the modular orbifold $\mathrm{PSL}_2\mathbb{Z} \setminus \mathbb{H}^2$ is well-known to be homeomorphic to both the quotient space $\mathrm{PSL}_2\mathbb{Z} \setminus \mathrm{PSL}_2\mathbb{R} \cong \mathrm{SL}_2\mathbb{Z} \setminus \mathrm{SL}_2\mathbb{R}$ and the exterior of a trefoil K in S^3. A flow on $\mathrm{PSL}_2\mathbb{Z} \setminus \mathrm{PSL}_2\mathbb{R}$ historically called *the geodesic flow* is defined by multiplying $\left(\begin{smallmatrix} e^t & 0 \\ 0 & e^{-t} \end{smallmatrix}\right)$ on the right, and its closed orbits are called *modular knots*. For each primitive hyperbolic element γ in $\mathrm{SL}_2\mathbb{Z}$, we may define the corresponding modular knot C_γ by $C_\gamma(t) = M_\gamma \left(\begin{smallmatrix} e^t & 0 \\ 0 & e^{-t} \end{smallmatrix}\right)$ $(0 \leq 0 \leq \log \xi_\gamma)$, where $M_\gamma^{-1}\gamma M_\gamma = \left(\begin{smallmatrix} \xi_\gamma & 0 \\ 0 & \xi_\gamma^{-1} \end{smallmatrix}\right)$ with $\xi_\gamma > 1$. Every modular knot admits such a presentation. By virtue of Bonatti–Pinsky's nice compactification [BP20], we obtain the following:

Theorem 4 ([Uek21b, Theorem 4]). *Modular knots and the missing trefoil in S^3 obey the Chebotarev law, if ordered by length in a generic metric.*

As a corollary, we obtain a result on a function with arithmetic origin. The discriminant function $\Delta(z) = q\prod_{n=1}^{\infty}(1-q^n)^{24}$ with $q = e^{2\pi\sqrt{-1}z}$, $z \in \mathbb{H}^2$ is a well-known modular function of weight 12. *The Dedekind symbol* Φ and *the Rademacher symbol* Ψ are the functions $\mathrm{SL}_2\mathbb{Z} \to \mathbb{Z}$ satisfying

$$\log\Delta(\gamma z) - \log\Delta(z) = \begin{cases} 6\log(-(cz+d)^2) + 2\pi i\Phi(\gamma) & \text{if } c \neq 0, \\ 2\pi i\Phi(\gamma) & \text{if } c = 0, \end{cases}$$

$$\Psi(\gamma) = \Phi(\gamma) - 3\mathrm{sgn}(c(a+d))$$

for any $\gamma = \left(\begin{smallmatrix} a & b \\ c & d \end{smallmatrix}\right) \in \mathrm{SL}_2\mathbb{Z}$ acting on $z \in \mathbb{C}$ via the Möbius transformation $\gamma z = \frac{az+b}{cz+d}$. Here we take a branch of the logarithm so that $-\pi \leq \mathrm{Im}\log z < \pi$ holds. This Ψ factors through the conjugacy classes of $\mathrm{PSL}_2\mathbb{Z}$ and satisfies $\Psi(\gamma^{-1}) = -\Psi(\gamma)$ for any γ.

The Rademacher symbol Ψ is known to be a highly ubiquitous function. Indeed, Atiyah proved the equivalence of seven definitions rising from very distinct contexts [Ati87], whereas Ghys gave further characterizations ([BG92], [Ghy07, Sections 3.3–3.5], [DIT17, Appendix]), proving that *for each primitive hyperbolic $\gamma \in \mathrm{SL}_2\mathbb{Z}$, the linking number between the modular knot C_γ and the missing trefoil K coincides with the Rademacher symbol*, namely,

$$\mathrm{lk}(C_\gamma, K) = \Psi(\gamma)$$

holds. Theorem 4 for $\rho(\gamma) = \mathrm{lk}(C_\gamma, K) \bmod m$ together with some arguments yield the following.

Corollary 5 ([Uek21b, Corollary 9]). *Suppose that γ runs through primitive hyperbolic elements of $\mathrm{SL}_2\mathbb{Z}$. For any $m \in \mathbb{Z}_{>0}$ and $k \in \mathbb{Z}/m\mathbb{Z}$, we have*

$$\lim_{\nu\to\infty} \frac{\#\{\gamma \mid |\mathrm{tr}\gamma| < \nu,\ \Psi(\gamma) = k \text{ in } \mathbb{Z}/m\mathbb{Z}\}}{\#\{\gamma \mid |\mathrm{tr}\gamma| < \nu\}} = \frac{1}{m}.$$

The similar arguments may be applicable to other Fuchsian groups. Modular knots for triangle groups around any torus knot in S^3 will be finely studied in [MU21].

References

[Ati87] Michael Atiyah. The logarithm of the Dedekind η-function. *Math. Ann.*, 278(1-4):335–380, 1987.

[BG92] J. Barge and É. Ghys. Cocycles d'Euler et de Maslov. *Math. Ann.*, 294(2):235–265, 1992.

[BP20] Christian Bonatti and Tali Pinsky. Lorenz attractors and the modular surface. preprint. arXiv:2001.05733, January 2020.

[DIT17] W. Duke, Ö. Imamoḡlu, and Á. Tóth. Modular cocycles and linking numbers. *Duke Math. J.*, 166(6):1179–1210, 2017.

[Ghy07] Étienne Ghys. Knots and dynamics. In *International Congress of Mathematicians. Vol. I*, pages 247–277. Eur. Math. Soc., Zürich, 2007.

[Maz12] Barry Mazur. Primes, Knots and Po. Lecture notes for the conference "Geometry, Topology and Group Theory" in honor of the 80th birthday of Valentin Poenaru, July 2012.

[McM13] Curtis T. McMullen. Knots which behave like the prime numbers. *Compos. Math.*, 149(8):1235–1244, 2013.

[Mih19] Tomoki Mihara. Cohomological approach to class field theory in arithmetic topology. *Canad. J. Math.*, 71(4):891–935, 2019.

[Mor12] Masanori Morishita. *Knots and primes*. Universitext. Springer, London, 2012. An introduction to arithmetic topology.

[MU21] Toshiki Matsusaka and Jun Ueki. Triangle modular knots and the Rademacher symbols. in preparation, 2021.

[Nii14] Hirofumi Niibo. Idèlic class field theory for 3-manifolds. *Kyushu J. Math*, 68(2):421–436, 2014.

[NU19] Hirofumi Niibo and Jun Ueki. Idèlic class field theory for 3-manifolds and very admissible links. *Trans. Amer. Math. Soc.*, 371(12):8467–8488, 2019.

[PP90] William Parry and Mark Pollicott. Zeta functions and the periodic orbit structure of hyperbolic dynamics. *Astérisque*, (187-188):268, 1990.
[Uek21a] Jun Ueki. Chebotarev links are stably generic. *Bull. Lond. Math. Soc.*, 53(1):82–91, February 2021.
[Uek21b] Jun Ueki. Modular knots obey the chebotarev law. preprint, March 2021.

НТБ ОНАХТ

I. Streltsova *Projective invariants of linear planar 3-webs* **145**

Kh. Sukhorukova *Ultrametric spaces of $*$-measures* **147**

B. M Sultanov *Sweep of surfaces in Galilean space* **148**

Y. Teplitskaya *Maximal distance minimizers. Examples and properties* **149**

M. Tkachuk, S. Plaksa *Analog of Menchov-Trokhimchuk theorem for monogenic functions in three-dimensional commutative algebra* **151**

E. Tosunoğlu, M. E. Kansu *Onto the some dynamic applications via quaternions* **153**

R. M. Tudoran *On dynamical systems with a prescribed globally bp-attracting set* **154**

T. P. Mokritskaya, A. V. Tushev *On certain fractal-based estimations of subsidence volume* **155**

Jun Ueki *Modular knots obey the Chebotarev law* **156**

J. Vielma, A. Guale *The Terence Tao set and the Collatz conjecture* **159**

V. Dilnyi, A. Vinskovska *Conformal mappings in Hardy-type spaces* **160**

I. Vlasenko *On the equivalency classes of weakly conjugated inner mappings* **161**

L. Vyhivska *Extremal problem for non-overlapping domains with free poles* **162**

N. Zorii *Inner harmonic measure for the fractional Laplacian* **163**

П. Задерей, М. Гаєвський *Мультиплікатори в просторах Харді та пов'язаних з ними просторах* **164**

М. І. Піструіл, І. М. Курбатова *Про деякі закономірності квазі-геодезичних відображень узагальнено-рекурентних просторів* **166**

С. М. Покась, И. И. Белокобыльский *Геодезические отображения римановых пространств второго приближения* **167**

НТБ ОНАХТ